



---

Sellan, G, Thompson, J, Majalap, N and Brearley, FQ (2019) Soil characteristics influence species composition and forest structure differentially among tree size classes in a Bornean heath forest. *Plant and Soil*, 438 (1-2). pp. 173-185. ISSN 0032-079X

---

**Downloaded from:** <https://e-space.mmu.ac.uk/622979/>

**Version:** Accepted Version

**Publisher:** Springer Verlag

**DOI:** <https://doi.org/10.1007/s11104-019-04000-5>

Please cite the published version

<https://e-space.mmu.ac.uk>

**Soil characteristics influence species composition and forest structure  
differentially among tree size classes in a Bornean heath forest.**

**Giacomo Sellan<sup>a</sup>, Jill Thompson<sup>b</sup>, Noreen Majalap<sup>c</sup> & Francis Q. Brearley<sup>a</sup>**

a) School of Science and the Environment, Manchester Metropolitan University, Chester  
Street, Manchester, M1 5GD, UK

b) Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB,  
Scotland, UK

c) Forest Research Centre, Sabah Forestry Department, PO Box 1407, 90715, Sandakan,  
Sabah, Malaysia

Corresponding author: [giacomo.sellan1@gmail.com](mailto:giacomo.sellan1@gmail.com)

**Cite as:** Sellan, G., Thompson, J., Majalap, N. Brearley, F.Q. Plant Soil (2019).

<https://doi.org/10.1007/s11104-019-04000-5>

**Abstract:**

*Background and aims:* Whilst several studies have shown that edaphic variability influences species composition in nutrient-poor tropical forests, the determinants of local species distributions and, in particular, how these change from younger to mature individuals in such forests are still under debate, and have been poorly explored in tropical heath forests that are among the least fertile tropical forest ecosystems.

*Methods:* We investigated the influence of soil fertility and topography on a Bornean heath forest species composition,  $\alpha$ -,  $\beta$ -diversity and tree size structure among size classes by recording all trees  $\geq 1$  cm DBH in 16 forest plots totalling 0.36 ha.

*Results:* Tree species distributions generally followed gradients in available Al and soil depth;  $\alpha$ - and  $\beta$ -diversity were linked to soil depth, and to some extent also to pH and the H:Al ratio. In contrast, forest structural attributes (basal area and stem density) were negatively correlated with both available and total P and a wider suite of soil nutrients, although trees  $\geq 10$  cm DBH were positively correlated with total P.

*Conclusion:* Our study shows that heath forest species distribution, richness and structure is related to both edaphic and topographic characteristics and that soil acidity might have a strong influence in shaping these forests' features. Among size classes, small trees are less influenced by soil and topography, whereas the sensitivity to these variables increases with tree size. We thus highlight that multiple edaphic factors influence different aspects of tropical forest structure, including different tree life stages, and species composition.

**Keywords:** Al toxicity; Kabili-Sepilok Forest Reserve; kerangas; nutrient limitation; species diversity; spodosol; white sand forest.

**INTRODUCTION:**

Tropical forests have the highest tree species diversity in the world (Gentry 1988; Ghazoul and Sheil 2010; Whitmore 1990) but we still strive to understand how such diversity is created and maintained. Several theories have been proposed to address this question such as the Janzen–Connell hypothesis (Janzen 1970, Connell 1971), neutral theory (Hubbell 2001) and environmental filtering (Baldeck et al. 2013). Although the drivers explained by different hypotheses might operate simultaneously and might have divergent controlling factors among sites, environmental factors influence tree species distribution and community composition in a considerable number of studies (Wright 2002, Legendre et al. 2005). In particular, climatic factors are important at continental scales (ter Steege et al. 2006, Zhang et al. 2016), whereas at smaller scales, topography and soil chemistry (Clark et al. 1998, Phillips et al. 2003) have a greater influence on species distributions (Condit et al. 2013, Jucker et al. 2018).

Most broad-scale studies of tree distribution only consider trees with a stem diameter at breast height (1.3 m; DBH) greater than 5 cm or 10 cm (e.g. Slik et al. 2015), so younger trees with smaller DBH (i.e. saplings) are often overlooked. In Borneo, for example, Paoli et al. (2006) and Sukri et al. (2012) reported small DBH trees to be distributed irrespectively of soil nutrient concentrations, whereas the distributions of larger DBH individuals were significantly influenced by edaphic variables. This implies that through tree ontogeny, individuals become more susceptible to the selective pressure of soil fertility so only the species adapted to particular soil conditions survive and grow (Russo et al. 2005). It is noteworthy that many experimental studies of forest nutrient limitation focus on seedlings and saplings grown in pots with different nutrient additions (e.g. Brearley et al. 2007, Nilus et al. 2011). If selective *in situ* response to soil fertility differ among size classes, then conclusions from pot bioassays must be taken with caution. It is thus paramount to consider trees from smaller size classes when conducting *in situ* studies of soil influences on species distribution and forest structure.

Heath forests occur throughout the tropics on nutrient-poor sandy soil (podzols), with vegetation physiognomy characterised by a generally stunted appearance, sclerophyllous leaves and short, untapered stems (Richards 1936, Janzen 1974, Peace & Macdonald 1981, Turner 1994, Becker et al. 1999, Proctor 1999, Turner et al. 2000, Kenzo et al. 2014). These forests also have a high density of understorey trees, low species diversity (Anderson 1981, Frasier et al. 2008), and a high degree of endemism (Garcia 2016). Heath forest distributions and the characteristic features of their trees have been ascribed to low soil pH (Luizao et al. 2007), deficient soil nitrogen (Luizao 1994, Proctor 1999, Proctor et al. 1983, Brearley et al. 2011) or the interaction of these two factors (Luizao 1996). A number of studies (Brünig 1974, Newbery 1991, Newbery et al. 1986) classified several typologies of Bornean heath forests (known as *kerangas* locally) but few have linked heath forest species distribution to either soil chemical or topographic variables. Nonetheless, these valuable studies draw some notable conclusions. At Gunung Mulu (Sarawak, Malaysia), Newbery & Proctor (1984) focussed on differences in soil chemistry and found species distribution to be associated with differences in soil organic C and CEC, whilst Miyamoto et al. (2003) found the most abundant tree species to be weakly associated with humus depth in a Central Kalimantan (Indonesia) heath forest. Din et al. (2015), instead, ascribed variation in a Bruneian heath forest community to soil N concentration but highlighted that other topographic and edaphic variables, not considered in their study, might also have an effect on species distributions.

In this study, we used sixteen small plots in heath forest situated in north-east Malaysian Borneo. All plots were in close proximity to one another (within an area c. 0.05 km<sup>2</sup>) to limit potential differences in bedrock and rainfall patterns and their influence on tree species distribution and forest structure. In particular, in this study we asked: (1) which soil chemical and topographic factors significantly influenced forest tree species composition and structure, and (2) how the influence of these factors varied among tree-size classes?

98

99 **METHODS:**

100 Our experimental site is the Kabili-Sepilok Forest Reserve (KSFR), located in the  
101 Malaysian state of Sabah on the island of Borneo (5° 51' N, 117° 58' E) (Figures 1a & 1b in  
102 Supplementary information). Local bedrock consists of sandstone interbedded with occasional  
103 mudstone inclusions. The climate is equatorial with an annual precipitation of c. 3000 mm;  
104 only one month (April) receives less than 100 mm on average (Fox 1973, Nilus 2003).

105 We installed sixteen 15 m x 15 m plots within the *kerangas* forest of the KSFR, all at  
106 least 30 m apart from one another (Figure 1c in Supplementary information) over podzol  
107 (USDA soil classification) soils. The plots were located on a gently sloping (c. 15 °), north-  
108 facing hillside. Plot slope was determined as the mean value of four measurements of the angle  
109 from the top to the bottom of each plot using a hypsometer (Vertex IV, Haglöf, Långsele,  
110 Sweden). Plot elevation was determined with a GPS (Garmin Etrex 10, Garmin Ltd, Kansas  
111 City, KS, USA). Within all plots, trees and lianas  $\geq 1$  cm diameter at breast height (DBH;  
112 diameter measured at 1.3 m from the ground) were permanently tagged and their DBH recorded.  
113 Furthermore, we recorded the height of 25 randomly selected trees per plot spanning the tallest  
114 to the shortest tree with a hypsometer (Vertex IV, Haglöf, Långsele, Sweden) to relate the DBH  
115 to position in the canopy. Lianas were measured at 1.3 metres from their last rooting point. We  
116 then binned trees with DBH:  $\geq 1$ -<2 cm,  $\geq 2$ -<5 cm,  $\geq 5$ -<10 cm and  $\geq 10$  cm (from now on  
117 referred as <2, 2-5, 5-10, and >10 cm DBH). Species identification was carried out by staff  
118 from Kabili-Sepilok Forest Research Centre Herbarium.

119 SOIL SAMPLING AND SOIL CHEMICAL ANALYSIS- Within every plot, a single soil pit of  
120 approximately 30 cm x 30 cm was dug to compare soil depths across the plots. Each plot was  
121 divided into four subplots (7.5 m x 7.5 m) and a soil sample from the top 5 cm was collected

in each subplot following removal of roots and coarse undecomposed leaves by digging a hole of approximately 10 cm x 10 cm using a knife; we collected 64 samples from the 16 plots in total. Each soil sample was split in two subsamples. The first subsample (approximately 2 g) was added to 30 ml of 1 M KCl and shaken for one hour in the field, allowed to equilibrate in a refrigerator for 18 hours, then filtered and analysed for  $\text{NH}_4^+$  and  $\text{NO}_3^-$  on a segmented flow analyser (Astoria-Pacific A2, Clackamas, OR, USA). On the second subsample, we measured moisture content, pH, available and total nutrients, C and N as well as exchangeable acidity and Al. Soil moisture content was determined gravimetrically after drying 3 g of soil to a constant weight at 105°C. To measure pH, 5 g of fresh soil was shaken in 12.5 ml of distilled water overnight and pH recorded with a Corning 240 pH meter. We then oven dried at 50°C and ground the remaining soil for use in macro and micro-nutrient analysis. Samples (0.25 g) were microwave-digested (Mars Xpress 5, CEM Corporation, Matthews, NC, USA) for total Al and P analysis with a solution of 8 ml  $\text{HNO}_3$  and 2 ml deionised  $\text{H}_2\text{O}$ . Extractable soil P, Fe and Mn were extracted with a Mehlich I solution (2 g of soil shaken with 20 ml of Mehlich I solution for 5 minutes on a reciprocating shaker at 120 oscillations per minute ;Wuenschel et al. 2015). For soil exchangeable Ca, Na, Mg and K, we added 2 g of soil to 20 ml of 1 M ammonium acetate and shook for two hours using a reciprocating shaker at 120 oscillations per minute (modified from Van Reeuwijk 2002). We determined exchangeable acidity ( $\text{H}^+$  and Al) through titration of a 1 M KCl extraction (5 g soil and 50 ml KCl shaken for one hour) with 0.02 M NaOH and 0.01 % phenolphthalein (modified from Van Reeuwijk 2002). We also measured exchangeable Al on the 1 M KCl extracts. Cation and metal concentrations were quantified using an iCAP Duo 6300 inductively coupled plasma optical emission spectrometer (Thermo Scientific, Waltham, MA, USA). Cation exchange capacity was calculated as sum of bases and exchangeable acidity. Total C and N concentrations were determined by combusting 0.15 g of soil in a Leco TruSpec CN analyser (St Joseph, MI, USA).

STATISTICAL ANALYSIS- For the ordinations, we used the package *vegan* (Oksanen 2015) in R 3.5.1 (R Development Core Team 2009). We visualised soil differences among plots by performing a redundancy analysis (RDA; significance checked with 999 permutations) of soil chemistry variables constrained by topography where all the variables were centred and scaled. We determined collinearity and selected the main variables that accounted for edaphic variation amongst plots for the canonical correspondence (CCA) analysis with a principal component analysis (PCA) following the approach of Abdi & Williams (2010). This consisted of selecting the variables that exceeded the expected average contribution to the two first principal components. Selected variables were then used as explanatory variables alongside topographical data for CCA ordination of our tree community dataset.

We calculated  $\alpha$ -diversity with the Shannon-Wiener ( $H'$ ) index, whereas for  $\beta$ -diversity we calculated a matrix of total dissimilarity between plots with Jaccard dissimilarity index (function *beta.pair()* of the R package *betapart*; Baselga & Orme 2012). A preliminary exploration of community composition across our plots was carried out using a detrended correspondence analysis (DCA). We then identified associations of selected species to soil variables following Shenbrot et al. (1991) who considered the position of the species centroids with respect to the standard error of the CCA ordination main centroid. If a species' centroid was located within the triplot main centroid's standard deviation, the species was labelled as a "generalist", whereas species that had a positive or negative position along an axis were considered a "specialist" (Table S3). The criterion to select species were a) their abundance among the plots using Shannon-Wiener diversity index and b) their score on the first two CCA axes. We centred and scaled the chemical, topographic variables as well as the absolute species abundances before the CCA, down-weighted the importance of rare species with the "*downweight()*" function in R package *vegan* (Oksanen 2015) and visualised the results using a CCA triplot. The significance of edaphic and topographic variables was obtained with a



permutational ANOVA (999 permutations; Oksanen 2015) by terms that also report a *pseudo*  $F$  test (i.e. the ratio of constrained and unconstrained total inertia, each divided by their respective degrees of freedom). We performed a non- parametric Spearman correlation to determine correlations among  $\alpha$ -diversity, stem density and basal area with the same chemical and topographical variables used in the CCA, whereas we performed a Mantel test (9999 permutations) to assess whether  $\beta$ -diversity correlated with soil or topographic variables selected with PCA. We used a Mantel test because  $\beta$ -diversity is presented as a dissimilarity matrix. The CCA, Spearman correlations and Mantel test were performed with all trees combined and then re-run with the four tree size classes (<2, 2-5, 5-10 and >10 cm DBH).

## RESULTS:

The forest plots were characterized by a typical short-statured heath forest with a canopy height of approximately 28 m. In total, we tagged 3336 living individuals  $\geq 1$  cm DBH. The understorey was dense with a mean of 0.73 trees 1-5 cm DBH per m<sup>2</sup> and 0.19 trees  $\geq 5$  cm DBH per m<sup>2</sup>; lianas were scarce (a mean across 16 plots of 217 individuals per ha). Trees < 2 cm DBH had a mean height of 2.9 m ( $\pm 0.9$  SD), trees 2-5 cm DBH had a mean height of 5.9 m ( $\pm 1.7$ ), trees 5-10 cm DBH had a mean height of 11.4 m ( $\pm 2.8$ ) and trees > 10 cm DBH had a mean height of 21.0 m ( $\pm 5.3$ ). Throughout our plots, there was a mean of 199.7 ( $\pm 30.1$ ) stems plot<sup>-1</sup> whereas basal area had a mean of 36.3 ( $\pm 10.9$ ) m<sup>2</sup> ha<sup>-1</sup>. Relative basal area and stem density for the 20 most abundant species is shown in Table 1.

**TREE DIVERSITY** - In total we identified 2398 trees and shrubs to species level and 784 trees to genus, 12 to family with 142 not identified. We found 124 species within 48 families of which Myrtaceae (19 %) and Rubiaceae (14%) were most abundant. Myrtaceae had the greatest basal area (31 %), followed by Dipterocarpaceae (19 %), Clusiaceae (11 %),

Euphorbiaceae (10 %) and Sapotaceae (5 %). Alpha diversity ( $H'$ ) throughout the plots had a mean of 3.48 ( $\pm 0.23$  SD). The DCA (Figure S2, Supplementary information) revealed three main plot groupings that reflected plot position on the hillslope, with strong floristic similarity between plot I, J and L near the ridge crest, A, F, G, H, K and M in the middle of the hill and B, C, D, E, N, O and P at the base (Figure S1c, Supplementary information).

TOPOGRAPHY, SOIL CHEMISTRY AND PCA GRADIENT - The plots lay on a gently sloping hill with an elevation spanning 50 m between the highest and the lowest plot. Plots at the top of the slope had deeper (i.e. > 90 cm in plot I) podzolised soil than lower elevation plots (i.e. 24 cm in plot D). Generally, all plots were acidic ( $\text{pH} < 4$ ) with a high concentration of total Al (mean  $247 \pm 28$  SD  $\mu\text{g g}^{-1}$ ) and a very low CEC (mean  $3.75 \pm 0.3$   $\text{cmol}_\text{c} \text{kg}^{-1}$ ), due largely to the high exchangeable acidity. Soil C:N ratio was high (mean  $22.9 \pm 0.7$ ). Low soil N was reflected by low ammonium (mean  $8.5 \pm 1.1$   $\mu\text{g g}^{-1}$ ) and nitrate (mean  $0.59 \pm 0.1$   $\mu\text{g g}^{-1}$ ) concentrations (Table S1). The RDA first axis was characterised by a plot slope gradient (loading on the first axis: 0.99,  $p < 0.05$ ) showing that flatter plots to have less acidic pH, and lower exchangeable acidity and CEC. The second RDA axis was characterised by plot elevation (loading on the second axis: 0.99,  $p < 0.01$ ) and, to a lesser extent, by soil depth (loading on the second axis: 0.83,  $p < 0.05$ ) showing that higher elevation plots had deeper soil with high available aluminium, low H:Al ratio and total as well as available P (Figure 1). The PCA found seven variables exhibited collinearity (total Al, exchangeable acidity and available Fe, Na, K, Mg and Ca) so these were removed along with variables with non-significant loadings. The PCA first and second axes explained 40.8 % and 28.8 % of the variability in soil properties, respectively. The first axis was related to CEC, total and available P, pH and  $\text{NH}_4^+$  whereas the second axis was linked to the H:Al ratio, available Al and Mn and the C:N ratio. The variables that exceeded the expected average contribution to the two first principal components were CEC, total P, available Al and P, H:Al ratio and pH (Table S2,

Supplementary information); these were retained for the CCA, Spearman and Mantel analyses along with topographic variables (elevation, slope) and soil depth.

SPECIES DISTRIBUTION RELATED TO ENVIRONMENTAL VARIABLES - From the CCA, we can see that species followed the first two axes of the ordination in agreement with topography and soil chemistry (Figure 2). Among the different tree DBH classes, soil chemistry and topography together explained between 68 % and 76 % of the floristic variation. In particular, soil depth and available Al were important for most size classes of trees but were less important for trees > 10 cm DBH ( $X^2 = 0.24$ ,  $p < 0.1$ , and  $X^2 = 0.25$ ,  $p > 0.1$ , for available Al and soil depth, respectively). Floristic variation in trees 5 - 10 cm DBH was influenced by CEC ( $X^2 = 0.21$ ,  $p < 0.05$ ) and total soil P ( $X^2 = 0.23$ ,  $p < 0.05$ ; Table 2). From the CCA ordination, we therefore identified two soil gradients along which species appear to be distributed. The first was the H:Al ratio-soil depth gradient and the second was the CEC-pH gradient (Figure 2). Although the CCA triplot's first axis was mainly driven by the H:Al ratio and soil depth gradients, available Al achieved high loadings on both the first (0.44; Table 2) and second axes (0.44) stressing the importance that Al retains in influencing species distributions.

SPECIES EDAPHIC ASSOCIATIONS AND SOIL GRADIENTS - From the CCA analyses, 51 species retained both high importance for among-plot diversity and high scores on the first two ordination axes. Of these species, 19 were associated with a low H:Al ratio and deep soil, whereas no species was associated with shallow soil and a high H:Al ratio (CCA first axis, Table S3). Furthermore, we identified another 18 species as generalists, but their association changed among the DBH classes considered. The only species that maintained a generalist species distribution among all DBH classes was *Cotylolebum melanoxylon* (Dipterocarpaceae). *Garcinia gaudichaudii* (Clusiaceae), *Gaertnera junghuhniana* (Rubiaceae), *Dracaena elliptica* (Asparagaceae), *Syzygium caudatilimum* (Myrtaceae) and *Syzygium* sp. were generalists in at least two size classes (Table S3).

FOREST STRUCTURE, ALPHA- AND BETA- DIVERSITY - Soil heterogeneity clearly influence forest structure although the Spearman and Mantel tests, consistent with the CCA, showed different influences of soil and topographic variables among DBH classes (Tables 3 & 4). In particular, stem density and basal area decreased with high soil P. Density of all trees, and particularly those <2 cm DBH ( $p < 0.01$ ) were negatively correlated with available Mn and density and basal area of trees 5-10 cm DBH was negatively correlated with CEC, available Al and pH in addition to slope and soil depth (all  $p < 0.05$ ).

A similar suite of elements also influenced  $\alpha$ -diversity but correlations with soil chemistry and topography were mostly found in trees 5-10 cm DBH (CEC and pH:  $p < 0.01$ ; total P, slope and depth:  $p < 0.05$ ) whereas in trees > 10 cm DBH  $\alpha$ -diversity correlated only with available Al and pH ( $p < 0.05$ ).

The results of the Mantel test for  $\beta$ -diversity (Table 4) showed topography to be consistently significant among size classes, although at different degrees and with different variables. In particular, soil depth was most significant for trees 2-5 cm DBH ( $p < 0.01$ ) and 5-10 cm DBH ( $p < 0.001$ ) along with plot elevation for trees 5-10 cm DBH ( $p < 0.01$ ). Soil chemistry was only correlated with  $\beta$ -diversity for trees 2-5 cm DBH (H:Al ratio at  $p < 0.05$ ) and 5-10 cm DBH (pH and H:Al ratio at  $p < 0.05$  and  $p < 0.01$ ).

## DISCUSSION:

Nutrient-poor tropical forests support high tree species diversity but the relative influence of environmental factors on these forests' floristic variability still needs further research. Our study site showed a typical heath forest floristic diversity, forest structure, and soil chemistry. We showed that soil acidity, topography and nutrient limitation act in concert

to shape species distribution, structure and diversity of the Kabili-Sepilok heath forest ecosystem. The influences of these topo-edaphic factors change among size classes.

The heath forest in our study site had a basal area of trees  $> 5$  cm DBH ( $34.4 \text{ m}^2 \text{ ha}^{-1}$ ) and stem density of trees  $> 5$  cm DBH ( $1997 \text{ stems ha}^{-1}$ ), similar to other Bornean heath forests (e.g. Proctor et al. 1983; Davies & Becker 1991; Miyamoto et al. 2007). When compared to the adjacent lowland evergreen rain forest, stem density was approximately doubled ( $1002 \text{ stems ha}^{-1}$ ; Nilus 2003) whilst basal area was very similar ( $35 \text{ m}^2 \text{ ha}^{-1}$ ; Nilus 2003). The most abundant families were rather different to other Bornean tropical heath forests. In comparison to the extensive study undertaken by Newbery (1991) in Brunei and Sarawak heath forests, our site lacked Myrsinaceae, Annonaceae and Chrysobalanaceae. This might be due to the limited area of our study ( $0.36 \text{ ha}$ ), requiring caution when comparing with results from other studies because such a small area might not be sufficient to exhaustively capture forest structure and biodiversity indices that are sample size dependent (Condit et al. 1996).

The soil (podzol) underlying our plots had a greater concentration of available P and a lower concentration of Al compared with the heath forest plots (acrisol) of Nilus (2003) in the same KSFR, suggesting that the soils in our plots had likely gone through a longer or more intense podzolisation process. During podzolisation, the soil is weathered and clay is eluviated (Bravard & Righi 1989) so Al and Fe are deposited in deeper layers of the mineral soil (Andriesse 1975), which often forms an indurated hardpan (Andriesse 1968). Usually, in tropical soils, available P is bound to Fe and Al, but the low concentration of these cations in podzols increases P availability in the soil solution (as also seen by Medina & Cuevas 1989, Coomes 1997, Metali et al. 2015).

INFLUENCE OF SOIL AND TOPOGRAPHY ON FLORISTIC DISTRIBUTION, DIVERSITY AND FOREST STRUCTURE - Our results show that topography and soil chemistry have a strong

influence on floristic distribution. Generally, it appears that, within our heath forest, the variation in species composition is driven by both edaphic and topographic variables, in particular available Al concentration and soil depth. On the other hand, soil nutrients (CEC and total P) along with soil pH play a major role only for tree (5 - 10 cm DBH)  $\alpha$ -diversity, whereas an interaction of soil acidity, Al and topographic factors influences  $\beta$ -diversity. As Al is an important constituent of soil acidity, we argue that, in combination with its ratio with  $H^+$ , is important in shaping heath forest species composition as hypothesised by Proctor (1999). It is noteworthy that available Al covaries with soil depth and elevation and is directly linked to a decrease in the H:Al ratio. In Brazilian cerrado ecosystems, a similar pattern of increasing Al with elevation was observed to underpin species composition (Guidão et al. 2002, Abreu et al. 2012, Soares et al. 2015, Silva et al. 2016). Aluminium is abundant in clay-rich acidic soils and, together with  $H^+$ , is toxic for plants. Along a soil gradient with different clay content (e.g. ultisol to podzol) Al is likely to create a species distribution gradient based on species' Al tolerance (Kidd & Proctor, 2001). On the other hand, within our sandy podzol, the lack of Al-rich clay means low soil Al concentrations (as seen in the plots at the base of the slope). In such cases, a beneficial role of soil Al would be to reduce  $H^+$  toxicity through its buffering action (Proctor 1999, Luizao 1996, Luizao et al. 2007). Hydrogen is more detrimental to plants than Al - experimental Al addition, for instance, enhanced growth of temperate (Kinraide 1993, Kidd & Proctor 2000) and tropical plants (Osaki et al., 1997) adapted to an extremely acidic Al-poor soil. In some acidic and sandy soils, it could be argued that plants accumulate Al in leaves to buffer  $H^+$  toxicity through their litterfall with Al accumulators composing more than 30 % of the biomass in e.g. Brazilian cerrado (Goodland & Pollard 1973, Haridasan 1982, 2008). In our plots, *Gaertnera junghuhniana* (Rubiaceae) the most common species with a generalist distribution, is an Al accumulator (i.e. has a leaf Al concentration higher than 1000  $\mu g g^{-1}$ ; Sellan 2019). This further supports the importance of soil acidity, and Al, as a driver of

species distribution in our study site. On the other hand, considering the emphasis given to low N availability in some studies as a theory to explain heath forest characters (e.g. see Vernimmen et al. 2013), N was too low to significantly explain differences among plots and thus was excluded from the analysis post PCA variable selection.

Soil P (available and total) influenced species richness of our plots for trees 5 - 10 cm DBH and structure (i.e. basal area and stem density) of trees > 5 cm DBH. Surprisingly, plots with high soil P concentrations had lower basal area and stem density (although this was largely driven by smaller trees < 10 cm DBH). We would have expected the contrary as P is often considered to be limiting in lowland rain forest (Cleveland et al. 2011; LeBauer and Treseder 2008), and evidence suggests that it might also be limiting in heath forest (Dent et al., 2006). High available P concentrations are found in plots with low exchangeable Al (as discussed above) where soil acidity is likely dominated by H<sup>+</sup> rather than Al (Proctor 1999). So, the negative relationship among soil P and forest structure is possibly due to H<sup>+</sup> toxicity rather than to soil P *per se*.

Interestingly, soil depth is a strong selective force for the distribution of trees 5 - 10 cm DBH with species as *Madhuca pallida* (Sapotaceae), *Shorea multiflora* (Dipterocarpaceae) and *Gluta oba* (Anacardiaceae) associated with deeper soil. This result seems to agree with the hypothesis proposed by Newbery (1991), and supported by Grainger & Becker (2001), that heath forests are composed of two ecophysiological and structural guilds: one with dominant deep-rooted trees and one with small shallow-rooted trees. Alternatively, soil depth is possibly important because a deeper soil has a greater volume and thus holds more nutrients per area unit.

DIFFERENT INFLUENCES OF EDAPHIC AND TOPOGRAPHIC VARIABLES AMONG SIZE CLASSES - The diverse influence of nutrients and topography on tree species distribution, plot  $\alpha$ - and  $\beta$ -

diversity as well as community structure clearly changes among size classes. In particular, trees <5 cm DBH showed weak correlations among structure, floristic variation and diversity with edaphic and topographic variables when compared to individuals > 5 cm DBH. This differential effect among size classes was documented elsewhere in Asian forests (e.g. Paoli et al. 2008, Sukri et al. 2012, Xu et al. 2016, Yang et al. 2016) and we propose this dissimilarity to be the effect of light competition. Small understorey trees are primarily limited by light availability as they are growing below the forest canopy (Coomes & Allen 2007, Cai et al. 2008) and only secondarily from other resources. Despite heath forest's low leaf area index (Vernimmen et al. 2007) and high light availability at ground level (Richards 1936), leaves of understorey saplings (Cao 2000, Cao and Booth 2001) had adaptations to low light availability when compared to leaves of the same species growing in brighter environments. Based on our results, indeed, the smallest size classes had a mean height of 2.9 m (< 2 cm DBH) and 5.7 m (2-5 cm DBH), well below the canopy height of 28 m, whereas trees 5-10 cm DBH had a mean height of c. 11 m, which might be sufficient to free them from light limitation. In this case, nutrient availability would have a greater influence on the distribution of trees >5 cm DBH. The poor correlation of tree species distribution with topo-edaphic factors for trees > 10 cm DBH might be explained by the dynamic nature of soil chemistry. In both tropical (Bauters et al. 2017) and temperate (Mueller et al. 2012; Vesterdal et al. 2008) forest plantations, changes in topsoil nutrient content and acidity have been recorded after just a few decades. Tropical trees can be as old as 1000 years (Chambers et al. 1998; Kurokawa et al. 2003) and in heath forests trees, diameter growth is slow compared to lowland evergreen forest (Nilus 2003; Brearley et al. unpublished data). Although we have no information on tree age in our forest, it is likely that trees > 10 cm DBH developed in a soil with different nutrient characteristics from the ones we see today. Alternatively, the lack of correlation between trees > 10 cm DBH species distribution and topo-edaphic variables might be due to the random mortality of big trees in our plots. It is



also interesting to note that trees > 10 cm DBH show an opposite correlation of basal area with soil total P and of  $\alpha$ -diversity with soil pH if compared to trees 5-10 cm DBH. This might highlight different strategies to overcome soil acidity and absorb nutrients between different tree life stages.

SOIL-SPECIES ASSOCIATIONS - We acknowledge that our study plots are small but, comparing our results with existing literature, we found them to be consistent with other studies. For example, Baltzer et al. (2005) found *Shorea multiflora* to be a sandstone forest (humult ultisol) specialist in the same Kabili-Sepilok Forest Reserve. Our results thus agree with Baltzer et al. (2005) because sandstone soil is higher in Al than heath forest and we found *S. multiflora* to be more common in soil with low H:Al ratio, i.e. a soil with high available Al. The same is applicable to the results of Potts et al. (2002) who found *Pimelodendron griffithianum* (Euphorbiaceae) to be a generalist, *Vatica micrantha* (Dipterocarpaceae) and *Mangifera* sp. (Anacardiaceae) to favour Al- rich humult ultisols and two *Diospyros* (Ebenaceae) species to be specialists in udult ultisols (more nutrient rich soil). The dispersion of species around the ordination centroid in our analysis revealed that more than half of the species considered in this analysis followed a soil gradient (i.e. 16 species out of 23, considering all size classes grouped together).

The cumulative number of species following the soil depth- H:Al gradient through the split size classes (18 species) instead of the CEC-pH gradient (19 species) was very similar. However, the higher score of the first CCA axis gives us a further reasonable criterion to suggest a greater influence of the depth-H:Al gradient rather than the CEC-pH gradient on species distribution. A secondary result is that, among the heath forest generalist species, we found *Gaertnera junghuhniana*, which is an Al accumulator. This characteristic might allow

391 *G. junghuhniana* to succeed throughout our heath forest site because of its capacity of attenuate  
392  $H^+$  toxicity through Al-rich leaf litterfall.

393

394 Our study demonstrated that there are complex interactions among topography, nutrient  
395 limitation and soil acidity that influence different aspects of forest tree species distributions  
396 and forest structure in this Bornean heath forest. These variables showed a greater importance  
397 for trees 5- 10 cm DBH, confirming a recently developed hypothesis suggesting that  
398 environmental filtering is almost absent in the early stages of plant development and become  
399 cumulative with age (Jabot et al. 2008). Generally, we suggest soil acidity to be a factor of  
400 primary importance in shaping this heath forest's floristic variation and structure. Given the  
401 high conservation value of this fragile forest type (Oktavia et al. 2015; Whitmore 1984) we  
402 underscore its susceptibility to changes in soil pH and encourage careful management and  
403 protection of heath forests.

**Acknowledgements:** This research was funded by Manchester Metropolitan University's Environmental Science Research Centre. We would like to thank the Sabah Biodiversity Council, the Soil Chemistry section, Ecology section and Herbarium staff from Sabah Forest Research Centre and in particular the research assistants Juanis Runcin, Jemson Jumian, Postar Miun, Spincer Sitim, Lioba Sawadon and Yun Len Lee. We are indebted to David McKendry and Graham Tinsley for the assistance in the laboratory, Yannis Kougoulous for preparing the map and Liam Trethowan, Matteo Giacomazzo and anonymous reviewers for the valuable suggestions and discussions.

**Conflict of Interest:** The authors declare that they have no conflict of interest.

#### **Literature Cited:**

- Abdi H, Williams LJ (2010) Principal component analysis. *Wiley Interdiscip. Rev. Comput. Stat.* 2:433–459.
- Abreu MF, Pinto JRR, Maracahipes L, Gomes L, DeOliveira EA, Marimon BS, Marimon BHJ, DeFarias J, Lenza E (2012) Influence of edaphic variables on the floristic composition and structure of the tree-shrub vegetation in typical and rocky outcrop cerrado areas in Serra Negra, Goiás State, Brazil. *Brazilian J. Bot.* 35:259–272.
- Anderson BA (1981) White-sand vegetation of Brazilian Amazonia. *Biotropica* 13:199–210.
- Andriesse JP (1968) A study of the environment and characteristics of tropical podzols in Sarawak (east-Malaysia). *Geoderma* 2:201–227.
- Baltzer JL, Thomas SC, Nilus R, Burslem DFRP (2005) Edaphic specialization in tropical trees: physiological correlates and responses to reciprocal transplantation. *Ecology* 86:3063–3077.
- Baldeck CA, Harms KE, Yavitt JB, John R, Turner BL, Valencia R, Navarrete H, Bunyavejchewin S, Kiratiprayoon S, Yaacob A, Supardi MNN, Davies SJ, Hubbell SP, Chuyong GB, Kenfack D, Thomas DW, Dalling JW (2013) Habitat filtering across tree life stages in tropical forest communities. *Proc. R. Soc. B.*

- 429 280:20130548.
- 430 Banin L, Lewis SL, Lopez-Gonzalez G, Baker TR, Quesada CA, Chao K, Burslem DFRP, Nilus R, Abu K,  
 431 Keeling HC, Tan S, Davies SJ, Mendoza AM, Vasquez R, Lloyd J, Neill DA, Pitman N, Phillips OL (2014)  
 432 Tropical forest wood production: a cross-continental comparison. *J. Ecol.* 102:1025–1037.
- 433 Bauters M, Verbeeck H, Doetterl S, Ampoorter E, Baert G, Vermeir P, Verheyen K, Boeckx P (2016) Functional  
 434 composition of tree communities changed topsoil properties in an old experimental tropical plantation.  
 435 *Ecosystems* 20:861–871
- 436 Becker P, Davies SJ, Moxin M, Ismail MZH, Simanjuntak PM (1999) Leaf size distributions of understorey  
 437 plants in mixed dipterocarp and heath forests of Brunei. *J. Trop. Ecol.* 15:123–128.
- 438 Bravard S, Righi D (1989) Geochemical differences in an Oxisol-Spodosol toposequence of Amazonia, Brazil.  
 439 *Geoderma* 44:29–42.
- 440 Brearley FQ, Scholes J, Press M, Palfner G (2007) How does light and phosphorous fertilisation affect the growth  
 441 and ectomycorrhizal community of two contrasting dipterocarp species? *Plant. Ecol.* 192:237–249.
- 442 Brearley FQ, Fine PVA, Perreijn K (2011) Does nitrogen availability have greater control over the formation of  
 443 tropical heath forests than water stress? A hypothesis based on nitrogen isotope ratios. *Acta Amaz.* 41:589–  
 444 592.
- 445 Brünig EF (1974) Ecological Studies in the Kerangas Forests of Sarawak and Brunei. Borneo Literature Bureau.  
 446 Kuching
- 447 Cai Z, Poorter L, Han Q, Bongers F (2008) Effects of light and nutrients on seedlings of tropical *Bauhinia* lianas  
 448 and trees. *Tree Physiol.* 28:1277–1285.
- 449 Cao KF (2000) Leaf anatomy and chlorophyll content of 12 woody species in contrasting light conditions in a  
 450 Bornean heath forest. *Can. J. Bot.* 78:1245–1253.
- 451 Cao KF, Booth EW (2001) Leaf anatomical structure and photosynthetic induction for seedlings of five  
 452 dipterocarp species under contrasting light conditions in a Bornean heath forest. *J. Trop. Ecol.* 17:163–175.
- 453 Chambers JQ, Higuchi N, Schimel JP (1998) Ancient trees in Amazonia. *Nature* 391:135–136.
- 454 Clark DB, Clark DA, Read JM (1998) Edaphic variation and the mesoscale distribution of tree species in a

- 455 neotropical rain forest. *J. Ecol.* 86:101–112.
- 456 Condit R, Engelbrecht BMJ, Pino D, Pérez R, Turner BL (2013) Species distributions in response to individual  
 457 soil nutrients and seasonal drought across a community of tropical trees. *Proc. Natl. Acad. Sci. U. S. A.*  
 458 110:5064–5068.
- 459 Coomes DA, Allen RB (2007) Mortality and tree-size distributions in natural mixed-age forests. *J. Ecol.* 95:27–  
 460 40.
- 461 Coomes DA, Dalponte M, Jucker T, Asner GP, Banin LF, Burslem DFRP, Lewis SL, Nilus R, Phillips OL, Phua  
 462 M, Qie L (2017) Area-based vs tree-centric approaches to mapping forest carbon in Southeast Asian forests  
 463 from airborne laser scanning data. *Remote Sens. Environ.* 194:77–88.
- 464 Coomes DA (1997) Nutrient status of Amazonian caatinga forests in a seasonally dry area: nutrient fluxes in litter  
 465 fall and analyses of soils. *Can. J. For. Res.* 27:831–839.
- 466 Connell JH (1971) On the role of natural enemies in preventing competitive exclusion in some marine animals  
 467 and in rain forest trees. In: *Proceedings of the Advanced Study Institute on Dynamics of Numbers in*  
 468 *Populations*, Oosterbeek (ed. P.J. den Boer and G. R. Gradwell), pp. 298–312. Centre for Agricultural  
 469 Publishing and Documentation, Wageningen
- 470 Curtin D, Campbell CA, Jalil A (1998) Effects of acidity on mineralisation: pH dependence of organic matter  
 471 mineralization in weakly acidic soils. *Soil Biol. Biochem.* 30:57–64.
- 472 Davies SJ, Becker P (1996) Floristic composition and stand structure of mixed dipterocarp and heath forests in  
 473 Brunei Darussalam. *J. Trop. For. Sci.* 8:542–569.
- 474 Dent DH, Bagchi R, Robinson D, Majalap-Lee N, Burslem DFRP (2006) Nutrient fluxes via litterfall and leaf  
 475 litter decomposition vary across a gradient of soil nutrient supply in a lowland tropical rain forest. *Plant Soil*  
 476 288:197–215.
- 477 Din H, Metali F, Sukri RS (2015) Tree diversity and community composition of the Tutong white sands, Brunei  
 478 Darussalam: a rare tropical heath forest ecosystem. *Int. J. Ecol.* 2015:807876.
- 479 Dubroeuq D, Volkoff B (1998) From oxisols to spodosols and histosols: evolution of the soil mantles in the Rio  
 480 Negro basin (Amazonia). *Catena* 32:245–280.
- 481 Duivenvoorden JE (1995) Tree species composition and rain forest-environment relationships in the middle

- 482 Caqueta area, Colombia, NW Amazonia. *Vegetatio* 120:91–113.
- 483 Frasier CL, Albert VA, Struwe L (2008) Amazonian lowland, white sand areas as ancestral regions for South  
484 American biodiversity: biogeographic and phylogenetic patterns in *Potalia* (Angiospermae: Gentianaceae).  
485 *Org. Divers. Evol.* 8:44–57.
- 486 Fox JED (1973) A Handbook to Kabili-Sepilok Forest Reserve, Sabah Forest Record No. 9. Borneo Literature  
487 Bureau, Kuching.
- 488 Garcia R, Dexter KG, Pennington T (2016) Amazonian white-sand forests show strong floristic links with  
489 surrounding oligotrophic habitats and the Guiana Shield. *Biotropica* 48:47–57.
- 490 Gentry A (1988) Changes in plant community diversity and floristic composition on environmental and  
491 geographical gradients. *Ann. Missouri Bot. Gard.* 75:1–34.
- 492 Ghazoul J, Sheil D (2010) *Tropical Rain Forest Ecology, Diversity, and Conservation*, New York: Oxford  
493 University Press.
- 494 Goodland R, Pollard R (1973) The Brazilian cerrado vegetation: a fertility gradient. *J. Ecol.* 61:219–224.
- 495 Grainger J, Becker P (2001) Root architecture and root : shoot allocation of shrubs and saplings in a Bruneian  
496 heath forest. *Biotropica* 33:363–368.
- 497 Haridasan M (1982) Aluminium accumulation by some cerrado native species of central Brazil. *Plant Soil* 65:265–  
498 273.
- 499 Haridasan M (2008) Nutritional adaptations of native plants of the cerrado biome in acid soils. *Brazilian J. Plant*  
500 *Physiol.* 20:183–195.
- 501 Hubbell SP (2001) *The Unified Neutral Theory of Biodiversity and Biogeography*, Princeton: Princeton  
502 University Press.
- 503 Janzen DH (1970) Herbivores and the number of tree species in tropical forests. *Am. Nat.* 104:501–528.
- 504 Janzen DH (1974) Tropical blackwater rivers, animals, and mast fruiting by the Dipterocarpaceae. *Biotropica*  
505 6:69–103.
- 506 John R, Dalling JW, Harms KE, Yavitt JB, Stallard RF, Mirabello M, Hubbell SP, Valencia R, Navarrete H,  
507 Vallejo M, Foster RB (2007) Soil nutrients influence spatial distributions of tropical tree species. *Proc. Natl.*

- 508 Acad. Sci. 104:864–869.
- 509 Jucker T, Bongalov B, Burslem DFRP, Nilus R, Dalponte M, Lewis SL, Phillips OL, Qie L, Coomes DA (2018)
- 510 Topography shapes the structure, composition and function of tropical forest landscapes. Ecol. Lett.
- 511 21:989–1000.
- 512 Kenzo T, Furutani R, Hattori D, Tanaka S, Sakurai K, Ninomiya I, Kendawang JJ (2014) Aboveground and
- 513 belowground biomass in logged-over tropical rain forests under different soil conditions in Borneo. J. For.
- 514 Res. 20:197–205.
- 515 Kidd PS, Proctor J (2000) Effects of aluminium on the growth and mineral composition of *Betula pendula* Roth.
- 516 J. Exp. Bot. 51:1057–1066.
- 517 Kidd PS, Proctor J (2001) Why plants grow poorly on very acid soils: are ecologists missing the obvious? J. Exp.
- 518 Bot. 52:791–799.
- 519 Kinraide T (1993) Aluminum enhancement of plant growth in acid rooting media. A case of reciprocal alleviation
- 520 of toxicity by two toxic cations. Physiol. Plant. 88:619–625.
- 521 Kurokawa H, Yoshida T, Nakamura T, Lai J, Nakashizuka T (2003) The age of tropical rain-forest canopy species,
- 522 Borneo ironwood (*Eusideroxylon zwageri*), determined by  $^{14}\text{C}$  dating. J. Trop. Ecol. 19:1–7.
- 523 Lebrija-Trejos E, Perez-Garcia EA, Meave JA, Bongers F, Poorter L (2010) Functional traits and environmental
- 524 filtering drive community assembly in a species-rich tropical system. Ecology 91:386–398.
- 525 Legendre P, Borcard D, Peres-Neto PR (2005) Analyzing beta diversity: partitioning the spatial variation of
- 526 community composition data. Ecol. Monogr. 75:435–450.
- 527 Legendre P, Fortin M, Borcard D (2015) Should the Mantel test be used in spatial analysis? Methods Ecol. Evol.
- 528 6:1239–1247.
- 529 Luizao FJ (1996) Ecological studies in contrasting forest types in central Amazonia. PhD Thesis, University of
- 530 Stirling.
- 531 Luizao FJ, Luizao RC, Proctor J (2007) Soil acidity and nutrient deficiency in central Amazonian heath forest
- 532 soils. Plant Ecol. 192:209–224.
- 533 Luizao RCC (1994) Soil biological studies in contrasting types of vegetation in central Amazonian rain forest.
- 534 PhD Thesis, University of Stirling.

- 535 Medina E, Cuevas E (1989) Patterns of nutrient accumulation and release in Amazonian forests of the upper Rio  
536 Negro basin. In: Mineral nutrients in tropical forest and savanna ecosystems. (Ed. Proctor J) Blackwell  
537 Scientific Publications, Oxford. 217–240.
- 538 Metali F, Abu Salim K, Tennakoon K, Burslem DFRP (2015) Controls on foliar nutrient and aluminium  
539 concentrations in a tropical tree flora: phylogeny, soil chemistry and interactions among elements. New  
540 Phytol. 205:280–292.
- 541 Miyamoto K, Suzuki E, Kohyama T, Seino T, Mirmanto E, Simbolon H (2003) Habitat differentiation among tree  
542 species with small-scale variation of humus depth and topography in a tropical heath forest of Central  
543 Kalimantan, Indonesia. J. Trop. Ecol. 19:43–54.
- 544 Miyamoto K, Rahajoe J, Kohyama T, Mirmanto E (2007) Forest structure and primary productivity in a Bornean  
545 heath forest. Biotropica. 39:35–42.
- 546 Mueller KE, Eissenstat DM, Hobbie SE, Oleksyn J, Jagodzinski AM, Reich PB, Chadwick OA, Chorover J (2012)  
547 Tree species effects on coupled cycles of carbon, nitrogen, and acidity in mineral soils at a common garden  
548 experiment. Biogeochemistry 111:601–614.
- 549 Newbery DM (1991) Floristic variation within kerangas (heath) forest: re-evaluation of data from Sarawak and  
550 Brunei. Vegetatio 96:43–86.
- 551 Newbery DM, Renshaw E, Brunig EF (1986) Spatial pattern of trees in kerangas forest, Sarawak. Vegetatio  
552 15:773–781.
- 553 Newbery DM, Proctor J (1984) Ecological studies in four contrasting lowland rain forests in Gunung Mulu  
554 National Park, Sarawak: IV. Associations between tree distribution and soil factors. J. Ecol. 72:475–495.
- 555 Nilus R (2003) Effect of edaphic variation on forest structure, dynamics and regeneration in a lowland tropical  
556 rainforest in Borneo. PhD Thesis, University of Aberdeen.
- 557 Oksanen J (2015) Multivariate analysis of ecological communities in R: vegan tutorial. R package version 2.4-5.
- 558 Oktavia D, Setiadi Y, Hilwan I (2015) The comparison of soil properties in heath forest and post-tin mined land:  
559 basic for ecosystem restoration. Procedia Environ Sci 28:124–131
- 560 Osaki M, Watanabe T, Tadano T (1997) Beneficial effect of aluminum on growth of plants adapted to low pH  
561 soils. Soil Sci. Plant Nutr. 43:37–41.



- 562 Paoli GD, Curran LM, Slik JWF (2008) Soil nutrients affect spatial patterns of aboveground biomass and emergent  
563 tree density in southwestern Borneo. *Oecologia* 155:287–299.
- 564 Paoli GD, Curran LM, Zak DR (2006) Soil nutrients and beta diversity in the Bornean Dipterocarpaceae: Evidence  
565 for niche partitioning by tropical rain forest trees. *J. Ecol.* 94:157–170.
- 566 Peace WJH, Macdonald FD (1981) An investigation of the leaf anatomy, foliar mineral levels, and water relations  
567 of trees of a Sarawak forest. *Biotropica* 13:100–109.
- 568 Peña-Claros M, Poorter L, Alarcon A, Blate G, Choque U, Fredericksen TS, Justiniano MJ, Leano C, Licona JC,  
569 Pariona W, Putz FE, Quevedo L, Toledo M (2012) Soil effects on forest structure and diversity in a moist  
570 and a dry tropical forest. *Biotropica* 44:276–283.
- 571 Phillips OL, Vargas PN, Lorenzo A, Cruz AP, Chuspe M, Sánchez WG, Yli-halla M, Rose S (2003) Habitat  
572 association among Amazonian tree species : a landscape-scale approach. *J. Ecol.* 91:757–775.
- 573 Potts MD, Ashton PS, Kaufman LK, Plotkin JB (2002) Habitat patterns in tropical rain forests: a comparison of  
574 105 plots in Northwest Borneo. *Ecology* 83:2782–2797.
- 575 Proctor J (1999) Heath forests and acid soils. *Bot. J. Scotl.* 51:1–14.
- 576 Proctor J, Anderson JM, Chai P, Vallack HW (1983) Ecological studies in four contrasting lowland rain forests  
577 in Gunung Mulu National Park, Sarawak: I. Forest environment, structure and floristics. *J. Ecol.* 71:237–  
578 260.
- 579 Van Reeuwijk LP (2002) Procedures for Soil Analysis. 6th edition. International Soil Reference and Information  
580 Centre, Wageningen, The Netherlands.
- 581 Vesterdal L, Schmidt IK, Callesen I, Nilsson LO, Gundersen P (2008) Carbon and nitrogen in forest floor and  
582 mineral soil under six common European tree species. *For. Ecol. Manage.* 255:35–48.
- 583 Richards PW (1936) Ecological observations on the rainforest of Mount Dulit, Sarawak. Part III. *J. Ecol.* 24:340–  
584 360.
- 585 Ruggiero PGC, Batalha MA, Pivello VR, Meilleres ST (2002) Soil-vegetation relationships in cerrado (Brazilian  
586 savanna) and semideciduous forest, southeastern Brazil. *Plant Ecol.* 160:1–16.
- 587 Russo SE, Davies SJ, King DA Tan S (2005) Soil-related performance variation and distributions of tree species

- 588 in a Bornean rain forest. *J. Ecol.* 93:879–889.
- 589 Shenbrot GI, Rogovin KA, Surov AV (1991) Comparative analysis of spatial organization of desert lizard  
590 communities in middle Asia and Mexico. *Oikos* 61:157–168.
- 591 Silva AC, Silva JL, Souza AF (2016) Determinants of variation in heath vegetation structure on coastal dune fields  
592 in northeastern South America. *Brazilian J. Bot.* 39:605–612.
- 593 Slik JWF, Raes N, Aiba SI, Brearley FQ, Cannon CH, Meijaard E, Nagamasu H, Nilus R, Paoli G, Poulsen AD,  
594 Sheil D, Suzuki E, van Valkenburg JLCH, Webb CO, Wilkie P, Wulffraat S (2009) Environmental  
595 correlates for tropical tree diversity and distribution patterns in Borneo. *Divers. Distrib.* 15:523–532.
- 596 Soares MP, Reys P, Pifano DS, deSa JL, da Silva PO, Santos TM, Silva FG (2015) Relationship between edaphic  
597 factors and vegetation in savannas of the Brazilian midwest region. *Rev. Bras. Ciência do Solo* 39:821–829.
- 598 ter Steege H, Pitman NCA, Phillips OL, Chave J, Sabatier D, Duque A, Molino, JF, Prevost MF, Spichiger R,  
599 Castellanos H, von Hildebrand P, Vasquez R (2006) Continental-scale patterns of canopy tree composition  
600 and function across Amazonia. *Nature* 443:444–447.
- 601 Sukri RS, Wahab RA, Salim KA, Burslem DFRP (2012) Habitat associations and community structure of  
602 dipterocarps in response to environment and soil conditions in Brunei Darussalam, northwest Borneo.  
603 *Biotropica* 44:595–605.
- 604 Sellan G (2019) Ecological responses of a Bornean heath forest to experimental lime and nitrogen fertilisation.  
605 PhD Thesis. Manchester Metropolitan University.
- 606 Turner IM (1994) Sclerophylly: primarily protective? *Funct. Ecol.* 8:669–675.
- 607 Turner IM, Lucas PW, Becker P, Wong SC, Yong JWH, Choong MF, Tyree T (2000) Tree leaf form in Brunei:  
608 a heath forest and a mixed dipterocarp forest compared. *Biotropica* 32:53–61.
- 609 Turner BL, Brenes-Arguedas T, Condit R (2018) Pervasive phosphorus limitation of tree species but not  
610 communities in tropical forests. *Nature* 555:367–370.
- 611 Vernimmen RRE, Bruijnzeel LA, Proctor J, Verhoef HA, Klomp NS (2013) Does water stress, nutrient limitation,  
612 or H-toxicity explain the differential stature among heath forest types in Central Kalimantan, Indonesia?  
613 *Biogeochemistry* 113:385–408.

- 614 Vitousek PM (1984) Litterfall, nutrient cycling, and nutrient limitation in tropical forests. *Ecology* 65:285–298.
- 615 Vitousek PM, Sanford RL (1986) Nutrient cycling in moist tropical forest. *Annu. Rev. Ecol. Syst.* 17:137–167.
- 616 Whitmore TC (1990) *An Introduction to Tropical Rain Forests*, Oxford: Clarendon Press.
- 617 Wright JJ (2002) Plant diversity in tropical forests: a review of mechanisms of species coexistence. *Oecologia*  
618 130:1–14.
- 619 Wuenscher R, Unterfrauner H, Peticzka R, Zehetner F (2002) A comparison of 14 soil phosphorus extraction  
620 methods applied to 50 agricultural soils from Central Europe. *Plant. Soil. Environ.* 61:86–96.
- 621 Xu W, Hao M, Wang J, Zhang C, Zhao X, von Gadow K (2016) Soil elements influencing community structure  
622 in an old-growth forest in northeastern China. *Forests* 7:159.
- 623 Yang QS, Shen GC, Liu HM, Wang ZH, Ma ZP, Fang XF, Zhang J, Wang XH (2016) Detangling the effects of  
624 environmental filtering and dispersal limitation on aggregated distributions of tree and shrub species: life  
625 stage matters. *Plos One* 11:e0156326.
- 626 Zhang C, Li X, Chen L, Xie G, Liu C, Pei S (2016) Effects of topographical and edaphic factors on tree  
627 community structure and diversity of subtropical mountain forests in the lower Lancang river basin. *Forests*  
628 7:222.
- 629
- 630

FIGURE 1. Redundancy analysis (RDA) biplot with the main soil chemical variables (grey) constrained by plot topography (blue) in the heath forest of the Kabili-Sepilok Forest Reserve, Sabah, Malaysia. Colour of the name of plots reflects their grouping on the DCA ordination.





FIGURE 2. Canonical correspondence analysis (CCA) triplot with the tree species (represented as crosses) of all size classes in study plots (represented as letters) in the heath forest of the Kabili-Sepilok Forest Reserve, Sabah, Malaysia. The species whose names are written in full are the most abundant following the Shannon-Wiener diversity index and have a significant loading on the first two ordination axes. The red circle represents the standard deviation of the ordination centroid and the colour of the name of plot reflects their grouping on the DCA ordination.



*Table 1 Stem density and basal area of the 20 most common species in heath forest plots in the Kabili-Sepilok Forest Reserve, Sabah, Malaysia.*

Species	% of total stem density	% of total basal area
<i>Gaertnera junghuhniana</i> (Rubiaceae)	12.74	1.48
<i>Diospyros fusiformis</i> (Ebenaceae)	7.45	0.65
<i>Syzygium caudatilimbium</i> (Myrtaceae)	6.07	1.25
<i>Pimelodendron griffithianum</i> (Euphorbiaceae)	5.63	10.12
<i>Dracaena elliptica</i> (Asparagaceae)	5.38	0.58
<i>Cotylelobium melanoxyton</i> (Dipterocarpaceae)	4.78	12.71
<i>Garcinia bancana</i> (Clusiaceae)	3.35	6.84
<i>Cleistanthus gracilis</i> (Phyllanthaceae)	3.13	1.78
<i>Actinodaphne borneensis</i> (Lauraceae)	3.00	0.97
<i>Tristaniaopsis obovata</i> (Myrtaceae)	2.75	19.22
<i>Chionanthus pluriflorus</i> (Oleaceae)	2.63	0.77
<i>Ternstroemia aneura</i> (Pentaphylacaceae)	2.53	0.89
<i>Shorea multiflora</i> (Dipterocarpaceae)	2.44	3.21
<i>Myrsine</i> sp. (Primulaceae)	1.85	0.39
<i>Calophyllum</i> sp. (Clusiaceae)	1.75	1.37
<i>Palaquium rostratum</i> (Sapotaceae)	1.19	3.26
<i>Eurycoma longifolia</i> (Simaroubaceae)	1.00	0.17
<i>Anisophyllea disticha</i> (Anisophylleaceae)	0.90	0.04
<i>Madhuca pallida</i> (Sapotaceae)	0.84	2.45
<i>Hancea griffithiana</i> (Euphorbiaceae)	0.66	0.09
Cumulative total	67.7	65.1

TABLE 2. Scores of soil and topographic variables on the first CCA axes in four tree size classes heath forest plots in the Kabili-Sepilok Forest

Reserve, Sabah, Malaysia. Significant values ('\*\*\*' < 0.001, '\*\*' < 0.01, '\*' < 0.05, '+' < 0.1) are in bold.

Size class (cm)		Available Mn	CEC	Available Al	Total P	Available P	pH	H:Al ratio	Plot slope	Soil depth	Plot elevation	Number of individuals
All	Axis1 score	-0.24	0.07	<b>0.44*</b>	-0.20	-0.31	-0.02	-0.54	<b>-0.23<sup>+</sup></b>	<b>0.85*</b>	0.67	3194
	F-stat	0.99	1.08	2.19	1.43	1.27	0.85	1.07	1.68	2.51	0.89	
<2	Axis1 score	0.21	-0.01	<b>-0.43*</b>	0.25	0.34	-0.08	0.58	0.35	<b>-0.83*</b>	-0.67	1413
	F-stat	0.86	0.92	2.11	1.16	1.30	0.95	1.21	1.43	1.89	0.78	
2-5	Axis1 score	-0.25	0.13	<b>0.41<sup>+</sup></b>	-0.10	-0.22	-0.13	-0.52	-0.11	<b>0.92*</b>	0.67	1087
	F-stat	0.90	1.11	1.50	1.08	1.06	0.89	0.92	1.39	2.36	0.90	
5-10	Axis1 score	0.09	<b>-0.14*</b>	<b>-0.35*</b>	<b>0.07*</b>	0.15	0.06	0.41	<b>0.19<sup>+</sup></b>	<b>-0.86**</b>	<b>-0.63<sup>+</sup></b>	353
	F-stat	1.30	1.59	1.94	1.75	1.00	1.34	1.42	1.50	2.61	1.14	
>10	Axis1 score	-0.30	0.09	<b>0.53<sup>+</sup></b>	-0.19	-0.35	-0.08	-0.58	-0.06	0.83	0.64	341
	F-stat	1.05	1.06	1.69	1.05	0.80	0.71	0.86	0.97	1.74	0.73	

TABLE 3. Results of the Spearman correlation test among forest structure (i.e. basal area and stem density) and  $\alpha$ -diversity (calculated with Shannon-Wiener diversity index) with edaphic and topographic variables in heath forest plots in the Kabili-Sepilok Forest Reserve, Sabah, Malaysia. Significant values ('\*\*\*' < 0.001, '\*\*' < 0.01, '\*' < 0.05) are in bold.

Size classes (cm)	Variable	Available Mn	CEC	Available Al	Total P	Available P	pH	H:Al ratio	Plot slope	Soil depth	Plot elevation
All	Basal area	0.30	0.15	-0.05	<b>0.61*</b>	0.37	-0.19	0.33	0.10	-0.18	-0.25
	Stem density	<b>-0.70**</b>	-0.35	0.003	<b>-0.54*</b>	<b>-0.60*</b>	0.08	-0.28	-0.17	-0.01	0.42
	$\alpha$ -diversity	-0.34	-0.20	0.01	0.01	-0.19	-0.02	0.08	-0.02	-0.40	-0.02
<2	Basal area	-0.57	-0.21	0.18	-0.38	-0.42	0.01	-0.37	-0.15	0.01	0.42
	Stem density	<b>-0.64**</b>	-0.15	0.26	-0.37	-0.49	-0.06	-0.45	-0.08	0.14	0.46
	$\alpha$ -diversity	-0.31	-0.22	0.04	-0.06	-0.13	0.01	0.08	0.05	-0.38	-0.15
2-5	Basal area	-0.27	-0.19	-0.18	-0.25	-0.20	0.03	0.09	-0.11	-0.02	0.08
	Stem density	-0.33	-0.07	-0.12	-0.09	-0.14	-0.15	0.21	0.18	-0.15	0.01
	$\alpha$ -diversity	-0.36	-0.17	0.06	0.03	-0.06	0.03	0.06	0.10	-0.39	-0.08
5-10	Basal area	<b>-0.50*</b>	<b>-0.70**</b>	<b>-0.49*</b>	<b>-0.74***</b>	<b>-0.62**</b>	<b>0.53*</b>	-0.09	<b>-0.51*</b>	-0.40	0.26
	Stem density	-0.45	<b>-0.82***</b>	<b>-0.65**</b>	<b>-0.73**</b>	<b>-0.54*</b>	<b>0.69**</b>	0.11	<b>-0.51*</b>	<b>-0.58*</b>	0.10
	$\alpha$ -diversity	-0.47	<b>-0.69**</b>	-0.37	<b>-0.56*</b>	-0.46	<b>0.66**</b>	0.14	<b>-0.54*</b>	<b>-0.61*</b>	0.09
>10	Basal area	0.36	0.32	0.13	<b>0.69**</b>	0.45	-0.29	0.19	0.10	-0.05	-0.15
	Stem density	0.10	0.19	-0.11	0.12	0.06	-0.09	0.05	-0.06	0.19	0.17
	$\alpha$ -diversity	0.02	0.44	<b>0.49*</b>	0.28	-0.05	<b>-0.54*</b>	-0.02	0.03	0.30	0.16



*Table 4. Results of the Mantel test between  $\beta$ -diversity (calculated as total dissimilarity matrix with Jaccard index) and edaphic and topographic variables in the Kabili-Sepilok Forest Reserve, Sabah, Malaysia. Significant values ('\*\*\*' < 0.001, '\*\*' < 0.01, '\*' < 0.05) are in bold.*

Size class (cm)	Available Mn	CEC	Available Al	Total P	Available P	pH	H:Al ratio	Plot slope	Soil depth	Plot elevation
All	-0.08	-0.27	0.03	-0.10	-0.06	-0.03	0.25	0.22	0.47	<b>0.30*</b>
<2	-0.09	-0.23	0.01	-0.05	-0.04	-0.01	0.20	0.21	<b>0.39*</b>	<b>0.29*</b>
2-5	-0.09	-0.17	0.10	-0.05	-0.09	0.09	<b>0.34*</b>	<b>0.39*</b>	<b>0.48**</b>	0.23
5-10	-0.01	-0.03	0.14	-0.06	-0.09	<b>0.25*</b>	<b>0.32**</b>	<b>0.27*</b>	<b>0.45***</b>	<b>0.32**</b>
>10	-0.11	-0.12	0.06	-0.02	-0.07	-0.07	0.06	-0.09	0.38	<b>0.35*</b>

**SUPPORTING INFORMATION:**

*FIGURE S1. Location of the study site. A) The island of Borneo with the approximate position of the Kabili-Sepilok Forest Reserve (Sabah, Malaysia) highlighted in red. B) The Kabili-Sepilok Forest Reserve with the approximate location of the study plots highlighted and C) the study plot positions in the forest. Colour of the name of plots reflects their grouping on the DCA ordination.*




FIGURE S2. Detrended correspondence analysis (DCA) using the absolute density of the 126 species found in 16 heath forest plots in the Kabili-Sepilok Forest Reserve, Sabah, Malaysia. Colour of the name of plots reflects their grouping on the DCA ordination.




TABLE S1. Soil chemical and topographical characteristics of heath forest plots in the Kabili-Sepilok Forest Reserve, Sabah, Malaysia. Values of chemical variables are means of four measurements whereas topographic variables consist of only one measurement per plot (excepting slope).

Plot	pH (H <sub>2</sub> O)	Moisture Content (%)	Exchangeable		H:Al Ratio	Total		Available		
			Acidity	Aluminium		Al	P	P	Mn	Fe
			cmol <sub>c</sub> kg <sup>-1</sup>					µg g <sup>-1</sup>		
A	3.89	5.37	1.51	0.18	7.7	178.7	40.8	10.7	2.4	20.2
B	4.05	8.74	2.02	0.32	9.1	148.6	61.1	21.5	6.3	18.1
C	3.75	5.79	2.42	0.55	3.5	216.8	42.5	10.7	0.9	33.1
D	3.56	13.90	4.26	0.76	4.7	271.8	70.2	16.6	2.3	29.7
E	3.68	6.67	2.10	0.37	5.2	153.0	36.7	8.2	0.9	17.8
F	3.82	4.00	1.47	0.26	4.6	106.5	28.2	8.6	0.7	14.4
G	3.63	9.40	2.77	0.42	3.4	287.9	59.3	15.4	4.2	42.8
H	3.56	6.49	4.31	1.13	2.8	365.1	56.9	12.4	1.1	33.7
I	3.80	6.11	2.18	0.68	2.2	402.5	42.8	10.7	1.1	44.6
J	3.72	7.02	2.53	0.80	2.8	535.0	38.8	9.5	1.2	42.3
K	3.68	5.80	2.36	0.63	2.9	260.4	57.6	15.4	1.0	22.5
L	3.43	13.60	3.74	0.60	5.3	285.9	70.1	15.2	1.3	21.5
M	3.74	4.62	2.50	0.40	5.2	173.7	54.2	19.8	2.7	18.2
N	3.82	6.20	1.64	0.31	4.7	133.8	38.1	9.4	0.7	15.5
O	3.64	7.04	2.28	0.39	4.9	209.0	65.6	18.1	3.0	22.7
P	3.58	8.43	2.77	0.47	5.2	236.3	76.2	22.9	6.2	28.8

Table 1. Continued.

Plot	Exchangeable				CEC	NH <sub>4</sub> <sup>+</sup>	NO <sub>3</sub> <sup>-</sup>	C:N Ratio	Plot ASL	Plot Slope	Soil Depth
	Ca	K	Mg	Na							
	cmol <sub>c</sub> kg <sup>-1</sup>										
A	0.30	0.20	0.61	0.12	2.37	4.9	0.0	26.6	55	9.5	23
B	0.13	0.07	0.11	0.03	3.59	4.2	0.0	26.3	49	12.3	19
C	0.16	0.12	0.37	0.07	3.36	9.6	0.2	24.5	69	13.7	23
D	0.14	0.19	0.44	0.08	6.03	11.7	1.1	25.2	53	14.6	24
E	0.14	0.13	0.32	0.05	2.59	6.4	0.4	23.7	50	14.8	27
F	0.16	0.10	0.21	0.04	2.00	6.2	0.6	17.7	73	13.3	29
G	0.19	0.20	0.38	0.09	4.68	10.0	0.8	26.0	66	13.1	32
H	0.21	0.17	0.82	0.23	5.67	5.4	0.5	19.5	64	14.1	39
I	0.16	0.13	0.35	0.07	2.64	7.8	0.8	21.7	83	13.3	100
J	0.13	0.20	0.33	0.12	3.11	7.6	0.2	16.7	74	12.8	49
K	0.19	0.18	0.50	0.07	3.47	11.6	0.7	24.6	61	14.3	39
L	0.17	0.37	0.75	0.09	6.00	9.8	0.0	21.9	64	14.8	48
M	0.34	0.17	0.58	0.06	3.96	8.9	0.8	27.0	60	14.9	32
N	0.12	0.12	0.23	0.05	2.25	1.6	0.3	21.7	51	14.1	21
O	0.49	0.20	0.65	0.07	3.66	9.0	2.0	22.7	42	15.8	27
P	0.51	0.20	0.78	0.07	4.62	20.8	1.1	21.5	48	15.4	33

*TABLE S2. Contribution on principal component analysis (PCA) first and second axis of soil chemical parameters from heath forest plots in the Kabili-Sepilok Forest Reserve, Sabah, Malaysia. The variables that scored a contribution higher than the expected one for both dimensions are noted in bold.*

Variable	Contribution to PC1	Contribution to PC2
<b>Av.Al</b>	6.89	15.41
<b>Av.Mn</b>	6.79	14.90
<b>Av.P</b>	12.75	7.88
C:N	1.20	14.57
<b>CEC</b>	17.36	2.38
<b>H:Al</b>	0.11	24.56
Moisture	11.58	0.58
NH <sub>4</sub>	11.47	0.81
NO <sub>3</sub>	0.84	9.26
<b>pH</b>	11.54	7.39
<b>Tot.P</b>	19.45	2.23

TABLE S3. Association of the 52 most important tree species with the first (H:Al ratio-soil depth) or second (CEC-pH) CCA axes ordination of heath forest plots in the Kabili-Sepilok Forest Reserve, Sabah, Malaysia. Columns are divided per size-class and axis considered. The species whose centroid was located within the triplot main centroid standard deviation have been called “generalists” (g in the table), whereas the species that had a positive or negative correlation with the indicated axis have been designated with a “+” or a “-”, respectively. If a cell is empty it means that the individuals from that species in that size class were not important in determining inter-plot variation.

Species	All		<2 cm		2-5 cm		5-10 cm		>10 cm	
	Depth- H:Al	CEC- pH	Depth- H:Al	CEC- pH	Depth- H:Al	CEC- pH	Depth- H:Al	CEC- pH	Depth- H:Al	CEC- pH
<i>Anisophyllea disticha</i>			+		+					
<i>Barringtonia</i> sp.	+									
<i>Calophyllum</i> sp.							-		g	g
<i>Chionanthus pluriflorus</i>										+
<i>Cleistanthus gracilis</i>	+		+		+					
<i>Cotylelebum melanoxyton</i>	g	g	g	g	g	g	g	g	g	g
<i>Dacryodes</i> sp.		+		+						
<i>Diospyros fusiformis</i>			g	g						
<i>Diospyros</i> sp.			+							
<i>Dracaena elliptica</i>	g	g	g	g	g	g				
<i>Elaeocarpus</i> sp.			+							
<i>Eurycoma longifolia</i>							+			
<i>Gaertnera junghuhniana</i>	g	g	g	g	g	g		-		
<i>Garcinia bancana</i>							g	g		
<i>Garcinia gaudichaudii</i>							g	g	g	g
<i>Gluta oba</i>	+						+			
<i>Glochidion</i> sp.		-								
<i>Gnetum</i> sp.			g	g				+		
<i>Gonystylus</i> sp.					g	g				
<i>Hancea griffithiana</i>							+			
<i>Horsfieldia</i> sp.						+				
<i>Ixonanthes reticulata</i>			g	g				+		
<i>Ixora</i> sp.		-								
<i>Koompassia malaccensis</i>						-				
<i>Litsea cylindrocarpa</i>	g	g					g	g		
<i>Madhuca pallida</i>	+		+				+			+
<i>Mangifera</i> sp.	+						+		+	

<i>Memecylon</i> sp.			+				
<i>Mezzettia</i> sp.			+				
<i>Myristica</i> sp.				+			
<i>Myristica malaccensis</i>					+		
<i>Myrsine</i> sp.		-		-			-
<i>Palaquium rostratum</i>		+		+		+	+
<i>Parinari</i> sp.		+					
<i>Parishia insignis</i>					+		+
<i>Pimeleodendron griffithianum</i>						g	g
<i>Psydrax</i> sp.		-	+		+		
<i>Rothmannia</i> sp.	+		+				
<i>Santiria</i> sp.	g	g				+	
<i>Shorea falciferoides</i>				-			
<i>Shorea multiflora</i>	+					+	
<i>Stemonurus</i> sp.						+	
<i>Syzygium caudatilmbum</i>	g	g			g	g	g
<i>Syzygium</i> sp.	g	g			g	g	g
<i>Syzygium tawahense</i>						+	
<i>Ternstroemia aneura</i>					g	g	
<i>Timonius flavescens</i>			g	g		-	
<i>Tristaniopsis obovata</i>				-			g
<i>Urophyllum arboreum</i>	+						g
<i>Vatica micrantha</i>					+		
<i>Xanthophyllum flavescens</i>						+	
<i>Xylopia ferruginea</i>		-				-	